**Drug Class:** Microbicides



## **Drug Description**

Dextran sulfate, also known as DS, is a heparin-like long-chain polysaccharide containing approximately 17% sulfur and up to three sulfate groups per glucose molecule. [1] [2]

#### **HIV/AIDS-Related Uses**

Dextran sulfate, an investigational drug, was previously studied for use as a potent inhibitor of HIV-1. Toxic adverse effects, including profound thrombocytopenia, prevented further in vivo research.[3]

Recent Phase I/II clinical studies have confirmed the safety and acceptability of dextran sulfate as a vaginal microbicide.[4] Dextran sulfate is now being evaluated as a microbicide to prevent the sexual transmission of HIV-1.[5]

## Non-HIV/AIDS-Related Uses

Dextran sulfate is an anticoagulant and antihyperlipidemic agent.[6] Dextran sulfate has also been evaluated in vitro and in vivo as a microbicide inhibitor of herpes simplex virus (HSV)-1, HSV-2, and human papillomaviruses.[7]

### **Pharmacology**

Dextran sulfate and other sulfated polysaccharides are potent and selective inhibitors of HIV-1 CXCR4 isolates.[8] Experiments carried out with dextran sulfate samples of increasing molecular weight and degree of sulfation have shown that antiviral activity increases with increasing molecular weight and degree of sulfation.[9]

Sulfated carbohydrates like dextran sulfate exert their inhibitory HIV effects by interacting with the positively charged amino acids in the V3 loop of the viral envelope glycoprotein gp120.[10] This interaction prevents viral attachment to the cell surface and the interaction between gp120 and the CD4 receptor on T cells.[11] [12]

Binding of gp120 varies depending on the isolate and may reflect the type of saccharide residues and

the degree and positions of sulfation. Molar percent inhibition differences seen between various sulfated compounds suggest a specific or preferential interaction rather than simple, nonspecific, negatively-charged sulfate interactions with the V3 loop.[13]

Sulfated carbohydrates show activity against HIV mutants that have become resistant to nucleoside reverse transcriptase inhibitors such as zidovudine, and they appear to develop drug resistance very slowly.[14]

HIV resistance to polysulfated carbohydrates may be mapped to single amino acid changes, including those outside the V3 loop region. In one resistance study, a single mutation located in the V2 loop of gp120, K178E, induced a 48-fold change in sensitivity to dextran sulfate. A single amino acid change at position 322 in the V3 loop (K322Q) conferred 62-fold virus resistance to dextran sulfate. These changes did not alter HIV-1 sensitivity to CXCR4 ligands, the fusion inhibitor enfuvirtide, or zidovudine. A previously found dextran sulfate mutation, Q310H, did not confer a change in virus sensitivity. This suggests that O310H is not relevant for direct resistance but may be needed to facilitate the emergence of drug-resistant mutations.[15]

Dextran sulfate and other sulfated carbohydrates may inhibit HSV through competitive inhibition for binding to glycosaminoglycans on the host cell surface.[16]

### **Adverse Events/Toxicity**

Continuous IV infusion of dextran sulfate is toxic, producing profound but reversible thrombocytopenia and extensive but reversible alopecia. Because of its toxicity, IV dextran sulfate is no longer studied for use in treating HIV.[17]

Intravaginal gel use of the drug is associated with minor adverse effects in clinical trials. Increased thirst and unexplained intermenstrual bleeding were observed but were not considered severe.[18] [19] No evidence of systemic toxicity or genital irritation was associated with twice daily gel use in



## **Adverse Events/Toxicity (cont.)**

HIV infected and uninfected women.[20] In a four-week study of gel acceptability when applied twice-daily or just prior to sexual activity, nearly all women using dextran sulfate said they would continue using or recommend use of the gel. Side effects associated with twice-daily use included itching, irritation, bleeding, slippery feeling, and a cold feeling. Women also complained of the gel interfering with sex due to excess lubrication and the need to apply the gel too frequently.[21] No differences were noted between women using 1% dextran sulfate gel compared to those using 4% gel.[22]

## **Drug and Food Interactions**

Sulfated polysaccharides such as dextran sulfate may act synergistically with other anti-HIV drugs (e.g., zidovudine).[23]

#### **Clinical Trials**

For information on clinical trials that involve Dextran sulfate, visit the ClinicalTrials.gov web site at http://www.clinicaltrials.gov. In the Search box, enter: Dextran sulfate AND HIV Infections.

## **Dosing Information**

Mode of Delivery: Intravaginal.[24]

Dosage Form: Intravaginal gel administered twice daily or just prior to sexual activity.[25] [26] Clinical trials have evaluated concentrations of 1% or 4% gel.[27]

#### Chemistry

CAS Name: Dextran, hydrogen sulfate[28]

CAS Number: 9042-14-2[29]

Molecular formula: H2-O4-S.x-Unspecified[30]

Molecular composition: 17% to 20% sulfur

content[31]

Molecular weight: 4,000 to 500,000 daltons[32]

Physical Description: White powder from alcohol plus ether.[33]

Solubility: Freely soluble in water.[34]

#### **Other Names**

DS[35]

Dextran sulfate sodium[36]

Dextrin sulphate[37]

DxS[38]

Dextrin sulfate[39]

## **Further Reading**

Clotet-Codina I, Armand-Ugon M, Clotet B, Este JA. Mapping of polyanion drug-resistance to a single amino acid of the gp120 of human immunodeficiency virus type 1. 13th International AIDS Conference, Barcelona. 2002. Abstract WePeA5653.

Kensinger RD, Catalone BJ, Krebs FC, Wigdahl B, Schengrund CL. Novel polysulfated galactose-derivatized dendrimers as binding antagonists of human immunodeficiency virus type 1 infection. Antimicrob Agents Chemother. 2004 May;48(5):1614-23. PMID: 15105112

Low-Beer N, Jespers V, McCormack S, Gabe R, Nunn A, Chapman A, Van Damme L, Lacey CJN. A safety study of dextrin sulphate gel as a novel vaginal microbicide: data from HIV negative and positive women. 15th International AIDS Conference, Bangkok. 2004. Abstract WeOrD1317.

Neurath AR, Strick N, Li YY. Anti-HIV-1 activity of anionic polymers: a comparative study of candidate microbicides. BMC Infect Dis. 2002 Nov 21:2(1):27. PMID: 12445331

Piret J, Lamontagne J, Bestman-Smith J, Roy S, Gourde P, Desormeaux A, Omar RF, Juhasz J, Bergeron MG. In vitro and in vivo evaluations of sodium lauryl sulfate and dextran sulfate as microbicides against herpes simplex and human immunodeficiency viruses. J Clin Microbiol. 2000



## **Further Reading (cont.)**

Jan;38(1):110-9. PMID: 10618073

Witvrouw M, De Clercq E. Sulfated polysaccharides extracted from sea algae as potential antiviral drugs. Gen Pharmacol. 1997 Oct;29(4):497-511. Review. PMID: 9352294

### **Manufacturer Information**

Dextran sulfate
Ueno Fine Chemicals Industry Ltd
1133 Sixth Ave / 27th Floor
New York, NY 10036
(212) 935-2323

#### **For More Information**

Contact your doctor or an AIDSinfo Health Information Specialist:

- Via Phone: 1-800-448-0440 Monday Friday, 12:00 p.m. (Noon) 5:00 p.m. ET
- Via Live Help: http://aidsinfo.nih.gov/live\_help Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET

#### References

- 1. Merck Index 2001; p. 520
- 2. ChemIDplus Available at: http://chem.sis.nlm.nih.gov/chemidplus/chemidlite.jsp. Accessed 01/03/05.
- 3. Antimicrob Agents Chemother 1991 Dec;25(12):2544-50
- 4. Microbicides Conf 1st, 2004. Abstract 02092\_2.
- 5. Intl AIDS Conf 13th, 2002. Abstract WeOrD1317.
- 6. Merck Index 2001; p. 520
- 7. Antimicrob Agents Chemother 2001 Dec;45(12):3427-32
- 8. Antimicrob Agents Chemother 2004 May;48(5):1614-23
- 9. Gen Pharmacol 1997 Oct;29(4):497-511
- 10. Antimicrob Agents Chemother 2004 May;48(5):1614-23
- 11. Gen Pharmacol 1997 Oct;29(4):497-511



- 12. J Clin Microbiol 2000 Jan;38(1):110-9
- 13. Antimicrob Agents Chemother 2004 May;48(5):1614-23
- 14. Gen Pharmacol 1997 Oct;29(4):497-511
- 15. Intl AIDS Conf 15th, 2004. Abstract WePeA5653.
- 16. J Clin Microbiol 2000 Jan;38(1):110-9
- 17. Intl AIDS Conf 13th, 2002. Abstract WeOrD1317.
- 18. Intl AIDS Conf 13th, 2002. Abstract WeOrD1317.
- 19. Microbicides Conf 1st, 2004. Abstract 02002.
- 20. Microbicides Conf 1st, 2004. Abstract 02002.
- 21. Microbicides Conf 1st, 2004. Abstract 02014.
- 22. Intl AIDS Conf 13th, 2002. Abstract WeOrD1317.
- 23. Gen Pharmacol 1997 Oct;29(4):497-511
- 24. Microbicides Conf 1st, 2004. Abstract 02092\_2.
- 25. Microbicides Conf 1st, 2004. Abstract 02014.
- 26. Microbicides Conf 1st, 2004. Abstract 02002.
- 27. Intl AIDS Conf 13th, 2002. Abstract WeOrD1317.
- 28. MeSH Available at: http://www.nlm.nih.gov/cgi.mesh/2004/MB\_cgi. Accessed 01/03/05.
- $29. \ ChemIDplus Available \ at: http://chem.sis.nlm.nih.gov/chemidplus/chemidlite.jsp. \ Accessed \ 01/03/05.$
- 30. ChemIDplus Available at: http://chem.sis.nlm.nih.gov/chemidplus/chemidlite.jsp. Accessed 01/03/05.
- 31. ChemIDplus Available at: http://chem.sis.nlm.nih.gov/chemidplus/chemidlite.jsp. Accessed 01/03/05.
- 32. Merck Index 2001; p. 520
- 33. Merck Index 2001; p. 520
- 34. Merck Index 2001; p. 520
- 35. NIAID DAIDS Anti-HIV Compounds Database Available at: http://www.niaid.nih.gov/daids/dtpdb/000040.htm. Accessed 01/03/05.
- 36. ChemIDplus Available at: http://chem.sis.nlm.nih.gov/chemidplus/chemidlite.jsp. Accessed 01/03/05.
- 37. Microbicides Conf 1st, 2004. Abstract 02092\_2.
- 38. Antimicrob Agents Chemother 2004 May;48(5):1614-23
- 39. BMC Infect Dis 2002 Nov21;2(1):27